



# STIC Search Report

## Biotech-Chem Library

STIC Database Tracking Number: 108474

**TO:** Devesh Khare  
**Location:** cm1/8a13/8b19  
**Art Unit:** 1623  
**Tuesday, November 18, 2003**

**Case Serial Number:** 10/007489

**From:** Susan Hanley  
**Location:** Biotech-Chem Library  
**CM1 6B05**  
**Phone:** 305-4053

**susan.hanley@uspto.gov**

Search Notes

1084774 PWSR request

USPTO  
Intranet

Home Index Resources Contacts Services Search

## Scientific and Technical Information Center

NPL Virtual Library &gt; Request a Prior Art Search

Patents Home | Site Feedback

[NPL Home | STIC Catalog | Site Guide | EIC | Automation Training/TRPs | Contact Us | STIC Staff | FAQ | Firewall Authentication]

mgi



## Commercial Database Search Request

10/15/03

Search requests relating to published applications, patent families, and litigation may be submitted by filling out this form and clicking on "Send."

For all other search requests, fill out the form, print, and submit the printout with any attachments to the STIC facility serving your Technology Center.

## Tech Center:

TC 1600    TC 1700    TC 2100    TC 2600  
 TC 2800    TC 3600    TC 3700    Other

## Enter your Contact Information below:

## Name:

Devesh Khare

Employee Number: 77931 Phone:

605-1199

Art Unit or Office: 1623 Building &amp; Room Number:

8 A 13, Mail 8 B19

Enter the case serial number (Required): 10/007,489

If not related to a patent application, please enter NA here.

Class / Subclass(es) 536/25.34

Earliest Priority Filing Date: 09/14/1998

## Format preferred for results:

Paper    Diskette    E-mail

## Provide detailed information on your search topic:

- In your own words, describe in detail the concepts or subjects you want us to search.
- Include synonyms, keywords, and acronyms. Define terms that have special meanings.
- \*For Chemical Structure Searches Only\*  
Include the elected species or structures, keywords, synonyms, acronyms, and registry numbers

108474

- \*For Sequence Searches Only\*  
Include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.
- \*For Foreign Patent Family Searches Only\*  
Include the country name and patent number.
- Provide examples or give us relevant citations, authors, etc., if known.
- FAX or send the abstract, pertinent claims (not all of the claims), drawings, or chemical structures to your EIC or branch library.

**Enter your Search Topic Information below:**

Please search the following claims:

Claim 1: A method for generating phosphorothioate oligo mixtures comprising:

1) growing a single-stranded recombinant DNA phage in modified media that uses thio-phosphate as a source of phosphate

2) harvesting the single-stranded phage and purifying the DNA corresponding to the recombinant DNA insert

3) fragmentation of the insert DNA such that oligo mixtures spanning the entire length of the segment are generated

Claim 2: the method of claim 1 used to generate phosphorothioate ds DNA, ss DNA, and/or RNA by in vivo incorporation of thio-phosphate into nucleotide precursor pools.

Thank you.  
devesh khare

**Special Instructions and Other Comments:**

(For fastest service, let us know the best times to contact you, in case the searcher needs further clarification on your search.)

Press ALT + F, then P to print this screen for your own information.

007 / 489

=> file medline

FILE 'MEDLINE' ENTERED AT 14:47:28 ON 18 NOV 2003

FILE LAST UPDATED: 13 NOV 2003 (20031113/UP). FILE COVERS 1958 TO DATE.

On April 13, 2003, MEDLINE was reloaded. See HELP RLOAD for details.

MEDLINE thesauri in the /CN, /CT, and /MN fields incorporate the MeSH 2003 vocabulary. See <http://www.nlm.nih.gov/mesh/changes2003.html> for a description on changes.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 158

L55	546 SEA FILE=MEDLINE ABB=ON PLU=ON 10101-88-9 OR THIOPHOSPHORIC OR 13598-51-1
L56	3 SEA FILE=MEDLINE ABB=ON PLU=ON L55 AND (SSDNA OR SINGLE-STRAN D? OR SS DNA)
L57	2090 SEA FILE=MEDLINE ABB=ON PLU=ON ORGANOTHIOPHOSPHORUS COMPOUNDS /CT
L58	1 SEA FILE=MEDLINE ABB=ON PLU=ON L56 AND L57

=> file embase

FILE 'EMBASE' ENTERED AT 14:47:29 ON 18 NOV 2003

COPYRIGHT (C) 2003 Elsevier Inc. All rights reserved.

FILE COVERS 1974 TO 13 Nov 2003 (20031113/ED)

EMBASE has been reloaded. Enter HELP RLOAD for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d que 173

L64	962 SEA FILE=EMBASE ABB=ON PLU=ON 10101-88-9 OR THIOPHOSPHORIC OR 13598-51-1 OR THIOPHOSPHATE
L65	69 SEA FILE=EMBASE ABB=ON PLU=ON L64 AND (SSDNA OR SINGLE-STRAND ? OR SS DNA OR ?PHAGE OR PLASMID)
L66	38 SEA FILE=EMBASE ABB=ON PLU=ON L65 AND ?OLIGO?
L70	15 SEA FILE=EMBASE ABB=ON PLU=ON L66 AND (HIGH OR THIOPHOSPHATE OR PHOSPHOROTHIOATE OR EXTENDING)/TI
L71	3 SEA FILE=EMBASE ABB=ON PLU=ON L70 NOT (CHIRAL OR EFFECT OR VIRAL OR VIVO OR ANTIPARALLEL OR SFII OR MICE OR MACROPHAGE OR GENE OR CPG)/TI
L72	1 SEA FILE=EMBASE ABB=ON PLU=ON L66 AND EXTENDING/TI
L73	4 SEA FILE=EMBASE ABB=ON PLU=ON L71 OR L72

=> file hcplus

FILE 'HCPLUS' ENTERED AT 14:47:30 ON 18 NOV 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of ACS, is strictly prohibited.

FILE COVERS 1907 - 18 Nov 2003 VOL 139 ISS 21  
FILE LAST UPDATED: 17 Nov 2003 (20031117/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

&gt; d que 110

L1 ( 1813)SEA FILE=HCAPLUS ABB=ON PLU=ON PHOSPHOROTHIOATE OLIGONUCLEOTI  
 DES+PFT,NT/CT  
 L2 231 SEA FILE=HCAPLUS ABB=ON PLU=ON L1(L)PREP/RL  
 L4 5 SEA FILE=REGISTRY ABB=ON PLU=ON O3PS/MF  
 L5 8754 SEA FILE=REGISTRY ABB=ON PLU=ON "PHOSPHOROTHIOATE"  
 L6 448 SEA FILE=REGISTRY ABB=ON PLU=ON L5 AND M/ELS  
 L7 35 SEA FILE=REGISTRY ABB=ON PLU=ON L6 NOT C/ELS  
 L8 40 SEA FILE=REGISTRY ABB=ON PLU=ON L4 OR L7  
 L9 354 SEA FILE=HCAPLUS ABB=ON PLU=ON L8  
 L10 2 SEA FILE=HCAPLUS ABB=ON PLU=ON L9 AND L2

&gt; d que 119

L3 1813 SEA FILE=HCAPLUS ABB=ON PLU=ON PHOSPHOROTHIOATE OLIGONUCLEOTI  
 DES+PFT,NT/CT  
 L4 5 SEA FILE=REGISTRY ABB=ON PLU=ON O3PS/MF  
 L5 8754 SEA FILE=REGISTRY ABB=ON PLU=ON "PHOSPHOROTHIOATE"  
 L6 448 SEA FILE=REGISTRY ABB=ON PLU=ON L5 AND M/ELS  
 L7 35 SEA FILE=REGISTRY ABB=ON PLU=ON L6 NOT C/ELS  
 L8 40 SEA FILE=REGISTRY ABB=ON PLU=ON L4 OR L7  
 L9 354 SEA FILE=HCAPLUS ABB=ON PLU=ON L8  
 L16 222353 SEA FILE=HCAPLUS ABB=ON PLU=ON DNA+PFT/CT  
 L17 5268 SEA FILE=HCAPLUS ABB=ON PLU=ON L16(L)(SS OR SINGLE-STRAND?)  
 L18 24 SEA FILE=HCAPLUS ABB=ON PLU=ON L17 AND L3  
 L19 1 SEA FILE=HCAPLUS ABB=ON PLU=ON L9 AND L18

&gt; d que 123

L3 1813 SEA FILE=HCAPLUS ABB=ON PLU=ON PHOSPHOROTHIOATE OLIGONUCLEOTI  
 DES+PFT,NT/CT  
 L16 222353 SEA FILE=HCAPLUS ABB=ON PLU=ON DNA+PFT/CT  
 L17 5268 SEA FILE=HCAPLUS ABB=ON PLU=ON L16(L)(SS OR SINGLE-STRAND?)  
 L18 24 SEA FILE=HCAPLUS ABB=ON PLU=ON L17 AND L3  
 L20 486562 SEA FILE=HCAPLUS ABB=ON PLU=ON (MONOTHIO? OR PHOSPHOROTHIO?  
 OR THIO? OR PHOSPHOROMONOTHIO?)  
 L21 24 SEA FILE=HCAPLUS ABB=ON PLU=ON L20 AND L18  
 L22 3 SEA FILE=HCAPLUS ABB=ON PLU=ON L21 AND (PHAGE OR BACTERIOPHAG  
 E)  
 L23 2 SEA FILE=HCAPLUS ABB=ON PLU=ON L22 NOT CIRCULAR/TI

&gt; d que 125

L1 ( 1813)SEA FILE=HCAPLUS ABB=ON PLU=ON PHOSPHOROTHIOATE OLIGONUCLEOTI  
 DES+PFT,NT/CT  
 L2 231 SEA FILE=HCAPLUS ABB=ON PLU=ON L1(L)PREP/RL  
 L16 222353 SEA FILE=HCAPLUS ABB=ON PLU=ON DNA+PFT/CT  
 L17 5268 SEA FILE=HCAPLUS ABB=ON PLU=ON L16(L)(SS OR SINGLE-STRAND?)  
 L24 12 SEA FILE=HCAPLUS ABB=ON PLU=ON L2 AND (PHAGE OR BACTERIOPHAGE  
 )  
 L25 2 SEA FILE=HCAPLUS ABB=ON PLU=ON L24 AND L17

&gt; d que 141

L4 5 SEA FILE=REGISTRY ABB=ON PLU=ON O3PS/MF  
 L5 8754 SEA FILE=REGISTRY ABB=ON PLU=ON "PHOSPHOROTHIOATE"  
 L6 448 SEA FILE=REGISTRY ABB=ON PLU=ON L5 AND M/ELS  
 L7 35 SEA FILE=REGISTRY ABB=ON PLU=ON L6 NOT C/ELS  
 L8 40 SEA FILE=REGISTRY ABB=ON PLU=ON L4 OR L7  
 L9 354 SEA FILE=HCAPLUS ABB=ON PLU=ON L8  
 L32 354 SEA FILE=HCAPLUS ABB=ON PLU=ON L9 AND L5  
 L33 11 SEA FILE=HCAPLUS ABB=ON PLU=ON L32 AND SINGLE-STRAND?  
 L39 10 SEA FILE=HCAPLUS ABB=ON PLU=ON L33 AND ?OLIGO?  
 L40 10 SEA FILE=HCAPLUS ABB=ON PLU=ON L39 AND ?THIO?  
 L41 7 SEA FILE=HCAPLUS ABB=ON PLU=ON L40 NOT (GOLD OR DOUBLE OR  
 HAPLOTYPES)/TI

&gt; s 110 or 119 or 123 or 125 or 141

L74 12 L10 OR L19 OR L23 OR L25 OR L41

&gt; dup rem 158 173 174

FILE 'MEDLINE' ENTERED AT 14:47:58 ON 18 NOV 2003

FILE 'EMBASE' ENTERED AT 14:47:58 ON 18 NOV 2003  
COPYRIGHT (C) 2003 Elsevier Inc. All rights reserved.FILE 'HCAPLUS' ENTERED AT 14:47:58 ON 18 NOV 2003  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)  
PROCESSING COMPLETED FOR L58  
PROCESSING COMPLETED FOR L73  
PROCESSING COMPLETED FOR L74  
L75 17 DUP REM L58 L73 L74 (0 DUPLICATES REMOVED)  
ANSWER '1' FROM FILE MEDLINE  
ANSWERS '2-5' FROM FILE EMBASE  
ANSWERS '6-17' FROM FILE HCAPLUS

-&gt; d ibib abs ind 1-5

L75 ANSWER 1 OF 17 MEDLINE on STN  
 ACCESSION NUMBER: 85054878 MEDLINE  
 DOCUMENT NUMBER: 85054878 PubMed ID: 6094546  
 TITLE: Cleavage of phosphorothioate-substituted DNA by restriction endonucleases.  
 AUTHOR: Potter B V; Eckstein F  
 SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (1984 Nov 25) 259 (22)  
 14243-8.  
 Journal code: 2985121R. ISSN: 0021-9258.  
 PUB. COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 198412  
 ENTRY DATE: Entered STN: 19900320  
 Last Updated on STN: 19900320  
 Entered Medline: 19841227

AB M13 RF DNA was synthesized in vitro in the presence of various single deoxynucleoside 5'-O-(1-thiotriphosphate) phosphorothioate analogues, and the three other appropriate deoxynucleoside triphosphates using a M13 (-)-single-stranded template, *Escherichia coli* DNA polymerase I and T4 DNA ligase. The resulting DNAs contained various restriction endonuclease recognition sequences which had been modified at their cleavage points in the (-)-strand by phosphorothioate substitution. The behavior of the restriction enzymes *Ava*I, *Bam*HI, *Eco*RI, *Hind*III, and *Sal*I towards these substituted DNAs was investigated. *Eco*RI, *Bam*HI, and *Hind*III were found to cleave appropriate phosphorothioate-substituted DNA at a reduced rate compared to normal M13 RF DNA, and by a two-step process in which all of the DNA is converted to an isolable intermediate nicked molecule containing a specific discontinuity at the respective recognition site presumably in the (-)-strand. By contrast, *Sal*I cleaved substituted DNA effectively without the intermediacy of a nicked form. *Ava*I, however, is only capable of cleaving the unsubstituted (+)-strand in appropriately modified DNA.

CT Check Tags: Support, Non-U.S. Gov't  
 Bacteriophage phi X 174: GE, genetics  
 Base Sequence  
 Binding Sites  
 \*DNA Restriction Enzymes: ME, metabolism  
 \*DNA, Single-Stranded: AN, analysis  
 DNA, Viral: AN, analysis  
 Deoxyribonuclease *Bam*HI  
 Deoxyribonuclease *Eco*RI  
 Deoxyribonuclease *Hind*III  
 \*Organothiophosphorus Compounds: ME, metabolism  
 \*Thiophosphoric Acid Esters: ME, metabolism

CN 0 (DNA, Single-Stranded); 0 (DNA, Viral); 0 (Organothiophosphorus Compounds); 0 (Thiophosphoric Acid Esters); EC 3.1.21 (DNA Restriction Enzymes); EC 3.1.21.- (Deoxyribonuclease *Bam*HI); EC 3.1.21.- (Deoxyribonuclease *Eco*RI); EC 3.1.21.- (Deoxyribonuclease *Hind*III); EC 3.1.21.- (endodeoxyribonuclease *Ava*I); EC 3.1.21.- (endodeoxyribonuclease *Sal*I)

L75 ANSWER 2 OF 17 EMBASE COPYRIGHT 2003 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN

ACCESSION NUMBER: 96053473 EMBASE  
 DOCUMENT NUMBER: 1996053473  
 TITLE: Extending the chemistry that supports genetic information transfer in vivo: Phosphorothioate DNA,

AUTHOR: Thaler D.S.; Liu S.; Tomblin G.  
 CORPORATE SOURCE: DNA RMCP, Jefferson Cancer Center, Thomas Jefferson University, 233 South 10th Street, Philadelphia, PA 19107, United States  
 SOURCE: Proceedings of the National Academy of Sciences of the United States of America, (1996) 93/3 (1352-1356).  
 ISSN: 0027-8424 CODEN: PNASA6

COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 022 Human Genetics  
 029 Clinical Biochemistry

LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB DNA and RNA are the polynucleotides known to carry genetic information in life. Chemical variants of DNA and RNA backbones have been used in structure-function and biosynthesis studies in vitro, and in antisense pharmacology, where their properties of nuclease resistance and enhanced cellular uptake are important. This study addressed the question of whether the base(s) attached to artificial backbones encodes genetic information that can be transferred in vivo. Oligonucleotides containing chemical variants of DNA or RNA were used as primers for site-specific mutagenesis of bacteriophage f1. Progeny phage were scored both genetically and physically for the inheritance of information originally encoded by bases attached to the nonstandard backbones. Four artificial backbone chemistries were tested: phosphorothioate DNA, phosphorothioate RNA, 2'-O-methyl RNA and methylphosphonate DNA. All four were found capable of faithful information transfer from their attached bases when one or three artificial positions were flanked by normal DNA. Among oligonucleotides composed entirely of nonstandard backbones, only phosphorothioate DNA supported genetic information transfer in vivo.

CT Medical Descriptors:

- \*gene transfer
- \*nucleotide sequence
- article
- chemical structure
- dna replication
- dna synthesis
- genetic code
- molecular genetics
- priority journal
- site directed mutagenesis
- structure activity relation

Drug Descriptors:

- \*dna
- \*rna
- antisense oligonucleotide
- oligonucleotide
- phosphorothioic acid
- transfer rna

RN (dna) 9007-49-2; (rna) 63231-63-0; (phosphorothioic acid) 10101-88-9, 13598-51-1, 15181-41-6; (transfer rna) 9014-25-9

L75 ANSWER 3 OF 17 EMBASE COPYRIGHT 2003 ELSEVIER INC. ALL RIGHTS RESERVED.  
 on STN

ACCESSION NUMBER: 93173385 EMBASE

DOCUMENT NUMBER: 1993173385

TITLE: Site-directed mutagenesis of single-stranded and double-stranded DNA by phosphorothioate approach.

AUTHOR: Olsen D.B.; Sayers J.R.; Eckstein F.  
 SOURCE: Methods in Enzymology, (1993) 217/- (189-217).  
 ISSN: 0076-6879 CODEN: MENZAU

COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 029 Clinical Biochemistry

LANGUAGE: English

CT Medical Descriptors:

- \*site directed mutagenesis
- article
- bacteriophage t7
- cell transformation
- dna sequence
- dna synthesis
- dna template

escherichia coli  
 gene mutation  
 hydrolysis  
 nonhuman  
 nucleotide sequence  
 plasmid  
 polymerization  
 priority journal  
 Drug Descriptors:  
 \*double stranded dna  
 \*phosphorothioic acid  
 \*plasmid dna  
 \*single stranded dna  
 dna polymerase  
 ethidium bromide  
 exodeoxyribonuclease iii  
 oligonucleotide  
 primer dna  
 restriction endonuclease  
 RN (phosphorothioic acid) 10101-88-9, 13598-51-1,  
 15181-41-6; (dna polymerase) 37217-33-7; (ethidium bromide) 1239-45-8;  
 (exodeoxyribonuclease iii) 9037-44-9

L75 ANSWER 4 OF 17 EMBASE COPYRIGHT 2003 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

ACCESSION NUMBER: 90080670 EMBASE  
 DOCUMENT NUMBER: 1990080670  
 TITLE: High-efficiency oligonucleotide  
 -directed plasmid mutagenesis.  
 AUTHOR: Olsen D.B.; Eckstein F.  
 CORPORATE SOURCE: Max-Plank Institut fur, Experimentelle Medizin, Abteilung  
 Chemie, Hermann-Rein Strasse 3, D-3400 Gottingen, Germany  
 SOURCE: Proceedings of the National Academy of Sciences of the  
 United States of America, (1990) 87/4 (1451-1455).  
 ISSN: 0027-8424 CODEN: PNASA6

COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 004 Microbiology  
 029 Clinical Biochemistry  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB A number of single- and double-base substitutions have been introduced into either the polylinker region or the lacZ gene in the plasmid vector pUC19. The efficiencies of these changes upon transfection of TG-1 bacterial cells were generally 70-80%. A strategy has been devised by which the wild-type DNA can be selectively destroyed. It is primarily based on the resistance of phosphorothioate internucleotide linkages to some restriction enzymes. A mismatch oligonucleotide is introduced into a gapped region and the gap is filled using three deoxynucleoside 5'-triphosphates and one deoxynucleoside 5'-[.alpha.-thio]triphosphate. Reaction with a restriction enzyme that is unable to hydrolyze phosphorothioates ensures that the DNA containing the mismatch oligonucleotide is only nicked. Concomitantly, the DNA that does not contain the desired mutation is linearized. Subsequent reactions with an exonuclease and DNA polymerase I yield mutant homoduplex DNA for transfection.

CT Medical Descriptors:  
 \*plasmid  
 \*site directed mutagenesis  
 genetic engineering  
 nonhuman  
 article  
 priority journal  
 Drug Descriptors:  
 \*phosphorothioic acid  
 RN (phosphorothioic acid) 10101-88-9, 13598-51-1,  
 15181-41-6

L75 ANSWER 5 OF 17 EMBASE COPYRIGHT 2003 ELSEVIER INC. ALL RIGHTS RESERVED.  
on STN

ACCESSION NUMBER: 90371788 EMBASE  
 DOCUMENT NUMBER: 1990371788  
 TITLE: Chemical and enzymatic ligation of 5'-thiophosphates of oligodeoxyribonucleotides  
 AUTHOR: Oshevskii S.I.  
 CORPORATE SOURCE: Institute of Cytology and Genetics, Siberian Branch of the  
 Academy of Sciences of the USSR, Novosibirsk, Russia

SOURCE: Doklady Biochemistry, (1990) 310/1-6 (15-18).  
 COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 029 Clinical Biochemistry  
 LANGUAGE: English  
 CT Medical Descriptors:  
 bacteriophage t4  
 article  
 Drug Descriptors:  
 \*dna  
 \*oligonucleotide  
 \*rna  
 RN (dna) 9007-49-2; (rna) 63231-63-0

=> d ibib abs hitrn 6

L75 ANSWER 6 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2003:461948 HCPLUS  
 DOCUMENT NUMBER: 139:225986  
 TITLE: Comparison of different antisense strategies in  
 mammalian cells using locked nucleic acids,  
 2'-O-methyl RNA, phosphorothioates and small  
 interfering RNA  
 AUTHOR(S): Grueneweller, Arnold; Wyszko, Eliza; Bieber, Birgit;  
 Jahn, Ricarda; Erdmann, Volker A.; Kurreck, Jens  
 CORPORATE SOURCE: Institut fuer Chemie-Biochemie, Freie Universitaet  
 Berlin, Berlin, D-14195, Germany  
 SOURCE: Nucleic Acids Research (2003), 31(12), 3185-3193  
 PUBLISHER: Oxford University Press  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Locked nucleic acids (LNAs) and double-stranded small interfering RNAs  
 (siRNAs) are rather new promising antisense mols. for cell culture and in  
 vivo applications. Here, we compare LNA-DNA-LNA gapmer  
 oligonucleotides and siRNAs with a phosphorothioate and  
 a chimeric 2'-O-Me RNA-DNA gapmer with respect to their capacities to  
 knock down the expression of the vanilloid receptor subtype 1 (VR1).  
 LNA-DNA-LNA gapmers with four or five LNAs on either side and a central  
 stretch of 10 or 8 DNA monomers in the center were found to be active  
 gapmers that inhibit gene expression. A comparative co-transfection study  
 showed that siRNA is the most potent inhibitor of VR1-green fluorescent  
 protein (GFP) expression. A specific inhibition was obsd. with an estd.  
 IC50 of 0.06 nM. An LNA gapmer was found to be the most efficient  
 single-stranded antisense oligonucleotide,  
 with an IC50 of 0.4 nM being 175-fold lower than that of commonly used  
 phosphorothioates (IC50 .apprx.70 nM). In contrast, the  
 efficiency of a 2'-O-methyl-modified oligonucleotide  
 (IC50.apprx.220 nM) was 3-fold lower compared with the  
 phosphorothioate. The high potency of siRNAs and chimeric LNA-DNA  
 oligonucleotides make them valuable candidates for cell culture  
 and in vivo applications targeting the VR1 mRNA.  
 IT 15181-41-6, Phosphorothioate  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (RNA; gene silencing using locked nucleic acids, 2'-O-Me RNA,  
 phosphorothioates and siRNA)  
 REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib abs hitrn 7

L75 ANSWER 7 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2003:339079 HCPLUS  
 DOCUMENT NUMBER: 139:1495  
 TITLE: Antisense technologies. Improvement through novel  
 chemical modifications  
 AUTHOR(S): Kurreck, Jens  
 CORPORATE SOURCE: Institut fur Chemie-Biochemie, Freie Universitat  
 Berlin, Berlin, 14195, Germany  
 SOURCE: European Journal of Biochemistry (2003) 270(8),  
 1628-1644  
 PUBLISHER: Blackwell Publishing Ltd.  
 DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review. Antisense agents are valuable tools to inhibit the expression of a target gene in a sequence-specific manner, and may be used for functional genomics, target validation and therapeutic purposes. Three types of anti-mRNA strategies can be distinguished. Firstly, the use of single stranded antisense-oligonucleotides; secondly, the triggering of RNA cleavage through catalytically active oligonucleotides referred to as ribozymes; and thirdly, RNA interference induced by small interfering RNA mols. Despite the seemingly simple idea to reduce translation by oligonucleotides complementary to an mRNA, several problems have to be overcome for successful application. Accessible sites of the target RNA for oligonucleotide binding have to be identified, antisense agents have to be protected against nucleolytic attack, and their cellular uptake and correct intracellular localization have to be achieved. Major disadvantages of commonly used phosphorothioate DNA oligonucleotides are their low affinity towards target RNA mols. and their toxic side-effects. Some of these problems have been solved in "second generation" nucleotides with alkyl modifications at the 2' position of the ribose. In recent years valuable progress has been achieved through the development of novel chem. modified nucleotides with improved properties such as enhanced serum stability, higher target affinity and low toxicity. In addn., RNA-cleaving ribozymes and deoxyribozymes, and the use of 21-mer double-stranded RNA mols. for RNA interference applications in mammalian cells offer highly efficient strategies to suppress the expression of a specific gene.

IT 15181-41-6, Phosphorothioate

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES

(Uses)

(comparison of different antisense strategy)

REFERENCE COUNT: 131 THERE ARE 131 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

&gt; d 1bib abs hitrn 8

L75 ANSWER 8 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN

ACCESSION NUMBER: 2003:609425 HCAPLUS

DOCUMENT NUMBER: 139:241236

TITLE: A comparison of gene repair strategies in cell culture

using a lacZ reporter system

AUTHOR(S): Nickerson, H. D.; Colledge, W. H.

CORPORATE SOURCE: Department of Physiology, University of Cambridge, Cambridge, UK

SOURCE: Gene Therapy (2003), 10(18), 1584-1591

CODEN: GETHEC, ISSN: 0969-7128

PUBLISHER: Nature Publishing Group

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Synthetic oligonucleotides and DNA fragments of less than 1 kilobase (kb) have been shown to cause site-specific genetic alterations in mammalian cells in culture and in vivo. We have used a lacZ reporter gene system to compare the efficiency of episomal and chromosomal gene repair in human embryonic kidney epithelial cells (HEK293), Chinese Hamster Ovary fibroblasts (CHOK1), human bronchial epithelial cells (16HBE), and mouse embryonic stem (ES) cells. The lacZ gene contains a G to A nucleotide change, (Glu to Lys mutation) that abrogates  $\beta$ -galactosidase activity. We compared the efficiency of different gene repair methods to correct this mutation and restore  $\beta$ -galactosidase activity. We evaluated PCR-generated double-stranded DNA fragments of 0.52-1.9 kb, single-stranded DNA oligonucleotides of 20, 35, or 80 bases contg. internal phosphorothioate links, and a 68 base RNA:DNA oligonucleotide. All of the oligonucleotides and DNA fragments showed some gene repair ability with an episomal plasmid. Short DNA fragments of 0.52 kb or greater gave the highest frequencies of episomal gene repair while single-stranded DNA oligonucleotides gave the highest frequency of chromosomal repair. In the context of a chromosomal target, antisense DNA oligonucleotides gave 5-fold higher frequencies of gene repair than their sense counterparts. The RNA:DNA chimeric oligonucleotide gave little or no gene repair on either a chromosomal or episomal target.

IT 15181-41-6, Phosphorothioate

RL: BSU (Biological study, unclassified); BUU (Biological use,

unclassified); BIOL (Biological study); USES (Uses)

(comparison of gene repair strategies in cell culture using a lacZ

reporter system)  
 REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib abs hitrn 9

L75 ANSWER 9 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:658292 HCAPLUS  
 DOCUMENT NUMBER: 137:196646  
 TITLE: Defined DNA sequences amplifiable with a universal primer pair for use in labeling materials for identification  
 INVENTOR(S): Brown, Tom; Thelwell, Nichola; Maxwell, Paula; Maxwell, Paul; Whiting, Paul  
 PATENT ASSIGNEE(S): Crime Solutions Limited, UK  
 SOURCE: PCT Int. Appl., 23 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002066678	A2	20020829	WO 2002-GB759	20020220
WO 2002066678	A3	20030530		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.: GB 2001-4163 A 20010220				
AB: A method of uniquely identifying an object by labeling it with a DNA sequence is described. The DNA sequence has a terminal region including a moiety that can be used to attach it to a substrate. Adjacent to this is a sequence by which the DNA can be released from the substrate, such as a restriction enzyme cleavage site. The remainder of the DNA is the unique identifier that includes a pair of primer binding sites sep'd. by a defined and unique DNA sequence. The DNA may also contain base analogs or have a modified backbone that will prevent degrdn. of the label by nucleases. The DNA may also be single-stranded with the immobilization region in the loop of a stem loop structure. The partially double stranded region may serve as a primer for an initial amplification. Amplification and sequencing of the unique sequence identifier can be used to demonstrate ownership.				
IT 15181-41-6D, Thiophosphate, nucleic acid conjugates RL: TEM (Technical or engineered material use); USES (Uses) (for immobilization of oligonucleotide label; defined DNA sequences amplifiable with universal primer pair for use in labeling materials for identification)				

=> d ibib abs hitrn 10

L75 ANSWER 10 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:575095 HCAPLUS  
 DOCUMENT NUMBER: 137:106042  
 TITLE: Nuclease-based method for detecting and quantitating oligonucleotides  
 INVENTOR(S): Yu, Zhengrong; Baker, Brenda F.; Wu, John  
 PATENT ASSIGNEE(S): Isis Pharmaceuticals, Inc., USA  
 SOURCE: PCT Int. Appl., 48 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002059137	A1	20020801	WO 2001-US49702	20011023
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				

CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
 GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT,  
 RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US,  
 UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,  
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF;  
 BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 EP 1337547 A1 20030827 EP 2001-994359 20011023  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR  
 PRIORITY APPLN. INFO.: US 2000-705587 A 20001103  
 WO 2001-US49702 W 20011023

AB The invention concerns a method for quantitating an oligonucleotide in a sample of bodily fluid and/or ext. is provided. The method comprises contacting an oligonucleotide with a probe comprising a detectable marker and a binding moiety; placing the fluid or ext. in contact with a solid support to which a binding partner of the binding moiety is attached; contacting the fluid or ext. with a single-strand specific nuclease to degrade probe which is not hybridized to the oligonucleotide; and detecting a label assocd. with the marker. The method provides for the detection and/or localization of oligonucleotides, including administered modified oligonucleotides, for therapeutic and/or pharmacokinetic purposes.

IT 15181-41-6, Phosphorothioate

RL: PRP (Properties)  
 (nuclease-based method for detecting and quantitating oligonucleotides)

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib abs hitrn 11

L75 ANSWER 11 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:522052 HCPLUS  
 DOCUMENT NUMBER: 137:89420  
 TITLE: Single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA)  
 INVENTOR(S): Bandaru, Rajanikanth; Kumar, Gyanendra  
 PATENT ASSIGNEE(S): Molecular Staging, Inc., USA  
 SOURCE: PCT Int. Appl., 90 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002053780	AZ	20020111	WO 2002-USS	20020104
WO 2002053780	A3	20030522		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2003044794	A1	20030306	US 2001-910372	20010720
US 6635425	B2	20031021		
EP 1347988	A2	20031001	EP 2002-705674	20020104
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 2003207323	A1	20031106	US 2003-465759	20030619
PRIORITY APPLN. INFO.:			US 2001-259918P	P 20010105
			US 2001-910372	A 20010720
			WO 2002-USS	W 20020104

AB The present invention provides a novel method for ligation of oligonucleotides contg. 5'-phosphorothioates on complementary templates by the action of DNA ligases. This reaction is readily applied to the synthesis of a single stranded circular DNA contg. a phosphorothioate. Directed ligation reaction by ATP dependent DNA ligase reaction is similar to conventional 5'-phosphate ligation. The utility of enzymic ligation in

probing specific sequences of DNA is also described. The present invention also provides a novel non-enzymic ligation of 5'-phosphorothioates that has been applied to the synthesis of single strand phosphorothioate and phosphate circular DNA. A process for detecting the presence of a mismatch in an otherwise complementary pair of oligonucleotides is disclosed using an enzyme-based technique which shows the presence of a mismatch by failing to form a ligated single stranded DNA circle that can optionally be amplified using std. methods of rolling circle amplification.

=> d ind 11

L75 ANSWER 11 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 IC ICM C12Q001-68  
 CC 3-1 (Biochemical Genetics)  
 Section cross-reference(s): 13  
 ST genotyping SNP single nucleotide polymorphism DNA high throughput assay; human genomic DNA SNP genotyping rolling circle amplification method; oligonucleotide rolling circle amplification nucleic acid  
 IT *Thermus thermophilus*  
 (DNA ligase from; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT *Escherichia coli*  
*Rhodothermus marinus*  
*Thermus scotoductus*  
 (DNA ligase; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT *Bacillus phage .phi.29*  
*Coliphage T4*  
*Coliphage T7*  
 (DNA polymerase; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Primers (nucleic acid)  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); PRP (Properties); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (DNA, Amplifluor, fluorescent labeled; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Genome  
 (DNA; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Alleles  
 (biallelic SNPs; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT RNA  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (bridging oligonucleotides contg.; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Peptides, biological studies  
 Primers (nucleic acid)  
 Proteins  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (closed circle oligonucleotides conjugates to; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Human  
 (genomic DNA polymorphisms; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Conformation  
 (hairpin loop, in oligonucleotide; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Enzymes, biological studies  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (mRNA-capping, single-stranded circular oligonucleotides synthesis using; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))  
 IT Class, uses

Plastics, uses  
 RL: DEV (Device component use); USES (Uses)  
 (oligonucleotide attached to solid support contg.; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Deoxyribonucleotides  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (open circle oligonucleotides and bridging oligonucleotides contg.; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT DNA  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); PRP (Properties); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (primer, Amplifluor, fluorescent labeled; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Nucleic acid amplification (method)  
 (rolling circle amplification; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Genetic polymorphism  
 (single nucleotide; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Genotyping (method)  
 Nucleic acid hybridization  
 (single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT DNA  
 RL: ANT (Analyte); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Probes (nucleic acid)  
 RL: ARG (Analytical reagent use); DGN (Diagnostic use); PRP (Properties); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (single-stranded circular, bridging, synthesis of; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Oligonucleotides  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)  
 (single-stranded circular, bridging, synthesis of; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT Phosphorothioate oligonucleotides  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); BIOL (Biological study); PREP (Preparation)  
 (single-stranded circular, synthesis of; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 9015-85-4, DNA ligase  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (E. coli, Thermus, Rhodothermus marinus, T4, single-stranded circular oligonucleotides synthesis using; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 9012-90-20, DNA polymerase, Klenow fragment  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (E. coli, phage T4 or T7, .phi.29, rolling circle amplification using; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 56-65-5, ATP, biological studies  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (as DNA ligase cofactor; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 7786-30-3, Magnesium chloride (MgCl<sub>2</sub>), biological studies  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (in ligation reaction buffer; single-stranded circular oligonucleotide

probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 25952-53-8, EDC  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (in single-stranded circular oligonucleotide synthesis; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 9037-46-1, Exonuclease I  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES  
 (Uses)  
 (ligation reaction products treated with; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 7704-34-9, Sulphur, biological studies  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (of 5'phosphorothioate group not used as bridging atom for single-stranded circular oligonucleotide synthesis; circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 9012-90-2, Taq DNA ligase 37259-52-2, Ampligase 37353-39-2  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); DGN (Diagnostic use); ANST (Analytical study); BIOL (Biological study); USES  
 (Uses)  
 (single-stranded circular oligonucleotides synthesis using; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

IT 440688-20-0 440688-21-1 440688-22-2 440688-23-3 440688-24-4, 5:  
 PN: WO02053780 SEQID: 5 unclaimed DNA 440688-25-5, 6: PN: WO02053780  
 SEQID: 6 unclaimed DNA 440688-26-6, 7: PN: WO02053780 SEQID: 7 unclaimed DNA 440688-27-7, 8: PN: WO02053780 SEQID: 8 unclaimed DNA 440688-28-8  
 440688-29-9 440688-30-2 440688-31-3 440688-32-4 440688-33-5  
 440688-34-6 440688-35-7 440688-36-8 440688-37-9 440688-38-0  
 440688-39-1 440688-40-4  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; single-stranded circular oligonucleotide probes for detection of polymorphisms in nucleic acids by rolling-circle amplification (RCA))

=> d ibib abs hitrn 12

L75 ANSWER 12 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:90226 HCAPLUS  
 DOCUMENT NUMBER: 136:145278  
 TITLE: Use of modified oligonucleotide to down-regulate gene expression  
 INVENTOR(S): Agrawal, Sudhir; Diasio, Robert B.; Zhang, Zhang  
 PATENT ASSIGNEE(S): Hybridon, Inc., USA  
 SOURCE: PCT Int. Appl., 71 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002008420	A2	20020231	WO 2001-US18338	20010606
WO 2002008420	A3	20021017		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, EG, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6608035	B1	20030819	US 2000-587934	20000606
PRIORITY APPLN. INFO.:			US 2000-587934	A 20000606
			US 1994-328520	A2 19941025
			US 1996-709910	B2 19960909
			US 1996-758005	B1 19961127

AB Disclosed is a method of down-regulating the expression of a gene in an animal, wherein a pharmacol. formulation comprising a chimeric oligonucleotide complementary to the gene is orally administered to an animal. The oligonucleotide administered has at least one

phosphorothioate internucleotide linkage and at least one alkylphosphonate, phosphorodithioate, alkylphosphonothioate, phosphoramidate, phosphoramidite, phosphate ester, carbamate, carbonate, phosphate triester, acetamide, or carboxymethyl ester internucleotide linkage.

IT 15181-41-6, Phosphorothioate  
 RL: ARC (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (internucleoside linkage; use of modified oligonucleotide to down-regulate gene expression)

=> d ind 12

L75 ANSWER 12 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 IC ICM C12N015-11  
 ICS C07H021-00; A61K031-7125; A61P025-28; A61P031-00; A61P033-00  
 CC 1-12 (Pharmacology)  
 Section cross-reference(s): 3, 14  
 ST modified oligonucleotide drug gene expression regulation  
 IT Lymphoma  
 (Burkitt's; use of modified oligonucleotide to down-regulate gene expression)  
 IT Trypanosoma cruzi  
 (Chagas' disease from; use of modified oligonucleotide to down-regulate gene expression)  
 IT Leukemia  
 (T-cell, adult; use of modified oligonucleotide to down-regulate gene expression)  
 IT Oligonucleotides  
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (acetamide linked; use of modified oligonucleotide to down-regulate gene expression)  
 IT Ameba  
 (amebiasis; use of modified oligonucleotide to down-regulate gene expression)  
 IT Gene  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (cellular, oligonucleotide is complementary to; use of modified oligonucleotide to down-regulate gene expression)  
 IT Disease, animal  
 (cryptosporidiosis, trichomoniasis; use of modified oligonucleotide to down-regulate gene expression)  
 IT Gene  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (expression; use of modified oligonucleotide to down-regulate gene expression)  
 IT Filaria  
 (filariasis; use of modified oligonucleotide to down-regulate gene expression)  
 IT Disease, animal  
 (foot-and-mouth disease; use of modified oligonucleotide to down-regulate gene expression)  
 IT Pathogen  
 Virus  
 (gene, oligonucleotide is complementary to; use of modified oligonucleotide to down-regulate gene expression)  
 IT Intestine, disease  
 (giardiasis; use of modified oligonucleotide to down-regulate gene expression)  
 IT Human herpesvirus 3  
 (herpes zoster from; use of modified oligonucleotide to down-regulate gene expression)  
 IT Ascarid  
 (infestation with, Ascariasis; use of modified oligonucleotide to down-regulate gene expression)  
 IT Pharynx, neoplasm  
 (nasopharynx, carcinoma; use of modified oligonucleotide to down-regulate gene expression)  
 IT Human herpesvirus  
 (oral and genital; use of modified oligonucleotide to down-regulate gene expression)  
 IT Drug delivery systems  
 (oral; use of modified oligonucleotide to down-regulate gene expression)  
 IT Wart

(papilloma; use of modified oligonucleotide to down-regulate gene expression)

IT Oligonucleotides  
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (phosphoramidite linked; use of modified oligonucleotide to down-regulate gene expression)

IT Schistosoma  
 (schistosomiasis from; use of modified oligonucleotide to down-regulate gene expression)

IT Toxoplasma gondii  
 (toxoplasmosis from; use of modified oligonucleotide to down-regulate gene expression)

IT AIDS (disease)  
 Alzheimer's disease  
 Blood plasma  
 Drug metabolism  
 Hepatitis  
 Influenza  
 Malaria  
 Mammalia  
 Parasite  
 Pneumocystis  
 Trichinella  
 Trichomonacides  
 (use of modified oligonucleotide to down-regulate gene expression)

IT Proteins  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (use of modified oligonucleotide to down-regulate gene expression)

IT Oligonucleotides  
 Phosphorothioate oligonucleotides  
 RL: BSU (Biological study, unclassified); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (use of modified oligonucleotide to down-regulate gene expression)

IT Human herpesvirus 3  
 (varicella from; use of modified oligonucleotide to down-regulate gene expression)

IT Papilloma  
 (warts; use of modified oligonucleotide to down-regulate gene expression)

IT Fever and Hyperthermia  
 (yellow; use of modified oligonucleotide to down-regulate gene expression)

IT 463-77-4, Carbamic acid, biological studies 993-13-5 3812-32-6, Carbonate, biological studies 7664-38-2D, Phosphoric acid, triesters, biological studies 13598-36-20, Phosphonic acid, alkyl 15181-41-6, Phosphorothioate 16481-04-2, Carboxy methyl ester 22638-09-1, Phosphoramidate  
 RL: ARG (Analytical reagent use); BSU (Biological study, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (internucleoside linkage; use of modified oligonucleotide to down-regulate gene expression)

IT 393599-15-0 393599-16-1 393599-17-2 393599-18-3 393599-19-4  
 393599-20-7 393599-21-8 393599-22-9 393599-23-0 393599-24-1  
 393599-25-2 393599-26-3 393599-27-4 393599-28-5 393599-29-6  
 RL: PRP (Properties)  
 (unclaimed nucleotide sequence; use of modified oligonucleotide to down-regulate gene expression)

&lt;- d ibib abs hitrn 13

L75 ANSWER 13 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2002:457395 HCPLUS  
 DOCUMENT NUMBER: 137:259481  
 TITLE: Separation of Synthetic Oligonucleotide Dithioates  
 from Monothiophosphate Impurities by Anion-Exchange  
 Chromatography on a Mono-Q Column  
 AUTHOR(S): Yang, Xianbin; Hodge, Richard P.; Luxon, Bruce A.;  
 Shope, Robert; Gorenstein, David G.  
 CORPORATE SOURCE: Sealy Center for Structural Biology and Department of  
 Human Biological Chemistry & Genetics, University of  
 Texas Medical Branch at Galveston, TX, 77555-1157, USA  
 SOURCE: Analytical Biochemistry (2002), 306(1), 92-99  
 CODEN: ANBCA2; ISSN: 0003-2697  
 PUBLISHER: Elsevier Science

DOCUMENT TYPE: Journal  
 LANGUAGE: English

A8 A method using a strong anion-exchange liq.-chromatog. column, Mono-Q, has been developed for high-resoln. anal. and purifn. of oligonucleotide dithioates, which were synthesized by an automated, solid-phase, phosphorothioamidite chem. High-resoln. sepn. of oligonucleotide phosphorodithioates from monothiophosphate impurities was obtained. High-resoln. sepn. was also demonstrated at pH 8. The sepn. of oligonucleotide dithioates was found to be linearly dependent on the no. of sulfurs for the same sequence length. Thiocyanate, SCN-, as eluting anion, can be used to purify oligonucleotides contg. a high percentage of phosphorodithioate linkages in lower salt concns. and provides better sepn. than chloride as eluting anion.

IT 15181-41-6P, Phosphorothioate  
 RL: BYP (Byproduct); PREP (Preparation)  
 (mono-, di-, sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)

REFERENCE COUNT: 39 THERE ARE 39 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ind 13

L75 ANSWER 13 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 CC 9-3 (Biochemical Methods)  
 Section cross-reference(s): 6  
 ST monoQ column oligonucleotide dithioate chromatog purifn; monothiophosphate oligonucleotide phosphorodithioate sepn  
 IT pH  
 (8, sepn. at; sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)  
 IT Ion exchange chromatography  
 (high-performance; sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)  
 IT Phosphorothioate oligonucleotides  
 RL: PUR (Purification or recovery); PREP (Preparation)  
 (sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)  
 IT 302-04-5, Thiocyanate, uses  
 RL: NUU (Other use, unclassified); USES (Uses)  
 (eluting anion of; sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange Chromatog. on a mono-Q Column)  
 IT 15181-41-6P, Phosphorothioate  
 RL: BYP (Byproduct); PREP (Preparation)  
 (mono-, di-, sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)  
 IT 131159-51-8, Mono Q HR 10/10  
 RL: NUU (Other use, unclassified); USES (Uses)  
 (sepn. of synthetic oligonucleotide dithioates from monothiophosphate impurities by anion-exchange chromatog. on a mono-Q column)

=> d ibib abs hitrn 14

L75 ANSWER 14 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 2001:868734 HCPLUS  
 DOCUMENT NUMBER: 136:1591  
 TITLE: Genotyping methods to detect DNA sequence polymorphisms and haplotypes  
 INVENTOR(S): Stanton, Vincent P., Jr.  
 PATENT ASSIGNEE(S): Variagenics, Inc., USA  
 SOURCE: PCT Int. Appl., 166 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001090419	A2	20011129	WO 2001-US16577	20010523
WO 2001090419	A3	20030710		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,

CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,  
 HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
 LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO,  
 RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ,  
 VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,  
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,  
 BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  
 US 6475736 B1 20021105 US 2000-696998 20001025  
 PRIORITY APPLN..INFO.: US 2000-206613P P 20000523  
 US 2000-696998 A2 20001025  
 US 2000-697013 A2 20001025  
 US 2000-697028 A2 20001025

AB Methods for detg. genotypes and haplotypes of genes are claimed. Also described are single nucleotide polymorphisms (SNPs) and haplotypes in the ApoE gene and their use in methods of this invention. Methods of the invention involve allele enrichment methods such as allele capture, allele-specific amplification, and allele-specific restriction endonuclease digestion. Allele capture means phys. sepn. of either single-stranded or double-stranded DNA. This can be accomplished by protein or nucleic acid reagents, such as disabled restriction enzymes, zinc-finger DNA-binding proteins, and covalent crosslinking agents, which have affinity for specific alleles. The captured complexes are then sepd. from the nucleic acid mixt. by reagents such as antibody-coated beads or streptavidin. Allele-specific amplification can be accomplished by strand obstruction, such as formation of stable secondary structures, or modified primers such as covalently crosslinkable primers. Lastly, allele-specific restriction methods for genotyping can be accomplished by triplex-mediated protection, primer-mediated creation of polymorphic restriction sites, and other variations, followed by amplification, direct nucleotide sequencing, or capture and size or sequence anal. Allele-specific primers were designed to det. haplotypes of nucleotide 186 T/C and 597 A/G polymorphisms in the dihydropyrimidine dehydrogenase gene. The primers are allele-specific because they induce hairpin loop formation when the "correct" nucleotide is present at the polymorphic site. The hairpin loop structure inhibits annealing of new primers and further amplification. PCR products were digested with BsrDI restriction endonuclease and analyzed by agarose gel electrophoresis. A T/C SNP at genomic site 21250 in the human ApoE gene results in a cysteine to arginine substitution at position 176 of the ApoE protein. For genotyping the T/C SNP, a loop primer and reverse primer were designed to amplify the target and introduce FokI and FspI restriction enzyme cleavage sites. Digestion with FokI and FspI produced allele-specific DNA fragments which were sequenced by mass spectrometry. Fourteen polymorphic sites for the ApoE gene and exptl. derived haplotypes for some or all of these polymorphisms are provided.

=> d ind 14

L75 ANSWER 14 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 IC ICM C120001-68  
 CC 3-1 (Biochemical Genetics)  
 Section cross-reference(s): 9, 13  
 ST genotyping polymorphism haplotype allele DNA binding complex restriction endonuclease; human gene ApoE SNP genotype haplotype PCR sequence analysis  
 IT Gene, animal  
 RL: ANT (Analyte); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (APOE; genotyping methods to detect DNA sequence polymorphisms and haplotypes)  
 IT Quaternary structure  
 (DNA triplex, allele-specific; genotyping methods to detect DNA sequence polymorphisms and haplotypes)  
 IT Ligands  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (DNA-binding; genotyping methods to detect DNA sequence polymorphisms and haplotypes)  
 IT Primers (nucleic acid)  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (DNA; genotyping methods to detect DNA sequence polymorphisms and haplotypes)  
 IT Enzymes, biological studies  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)  
 (RecA; genotyping methods to detect DNA sequence polymorphisms and

- IT      haplotypes)
- IT      Molecular association
  - (allele-specific DNA-binding; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Hydrogen bond
  - (allele-specific; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      RNA
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (aptamer; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Peptide nucleic acids
- IT      Proteins
- IT      Transcription factors
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (biotinylated or immobilized; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      DNA
  - RL: ANT (Analyte); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (double-stranded; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Alleles
- IT      Crosslinking
- IT      Genotypes
- IT      Genotyping (method)
- IT      Immunoassay
- IT      Nucleic acid amplification (method)
- IT      PCR (polymerase chain reaction)
- IT      RFLP (restriction fragment length polymorphism)
  - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Gene, animal
- IT      cDNA
  - RL: ANT (Analyte); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Oligonucleotides
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Peptide nucleic acids
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Phosphorothioate oligonucleotides
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Primers (nucleic acid)
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Proteins
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Transcription factors
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Peptides, biological studies
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (histidine-contg., ligand tag; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT      Oligonucleotides
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)

- (immobilized; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Oligonucleotides
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (labeled, biotinylated; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Magnetic particles
  - (ligand tag; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Antibodies
  - Avidins
    - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
      - (ligand tag; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Conformation
  - (loop, nucleic acid, D-loop, allele-specific; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT DNA sequence analysis
  - (mass spectrometric; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Nucleic acid bases
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
    - (mass-modified; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Imaging
  - (optical mapping; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Nucleic acid bases
  - RL: BSU (Biological study, unclassified); BIOL (Biological study)
    - (pairing, allele-specific; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT DNA
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (primer; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Genetic element
  - RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)
    - (restriction endonuclease cleavage site; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Polyamides, biological studies
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (sequence-specific DNA-binding; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Genetic polymorphism
  - (single nucleotide; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT DNA
  - RL: ANT (Analyte); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (single-stranded; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Separation
  - (size selection; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Immunoassay
  - (solid-phase; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Proteins
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (zinc finger-contg., biotinylated or immobilized; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT Proteins
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (zinc finger-contg.; genotyping methods to detect DNA sequence polymorphisms and haplotypes)
- IT 9012-90-2, DNA polymerase
  - RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST (Analytical study); BIOL (Biological study); USES (Uses)
    - (T4 and I, exonuclease; genotyping methods to detect DNA sequence polymorphisms and haplotypes)

IT 66-97-7, Psoralen 22542-10-5D, complexes, biological studies  
 146237-51-6 146237-52-7 146237-53-8  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST  
 (Analytical study); BIOL (Biological study); USES (Uses)  
 (crosslinking agent; genotyping methods to detect DNA sequence  
 polymorphisms and haplotypes)

IT 9026-89-5, Dihydropyrimidine dehydrogenase  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (gene for; genotyping methods to detect DNA sequence polymorphisms and  
 haplotypes)

IT 9037-44-9, Escherichia coli exonuclease III 9075-08-5, Restriction  
 endonuclease 37228-74-3, Exonuclease 37367-70-7, Lambda exonuclease  
 58513-62-5, Nuclease, bacteriophage T7 exodeoxyribo-  
 81295-34-3, Restriction endonuclease PvuII 81458-03-9, Restriction  
 endonuclease FokI 85340-94-9, Bal31 exonuclease 92228-44-9,  
 Restriction endonuclease NcoI 103780-20-7, NotI restriction endonuclease  
 107824-63-5 135340-89-5, Restriction endonuclease N.BstNBI  
 174632-11-2, Restriction endonuclease BsgI 189088-83-3, Restriction  
 endonuclease BsrDI  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST  
 (Analytical study); BIOL (Biological study); USES (Uses)  
 (genotyping methods to detect DNA sequence polymorphisms and  
 haplotypes)

IT 58-85-5, Biotin 7440-02-0, Nickel, biological studies 9013-20-1,  
 Streptavidin  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST  
 (Analytical study); BIOL (Biological study); USES (Uses)  
 (ligand tag; genotyping methods to detect DNA sequence polymorphisms  
 and haplotypes)

IT 9025-82-5, Phosphodiesterase  
 RL: ARG (Analytical reagent use); BUU (Biological use, unclassified); ANST  
 (Analytical study); BIOL (Biological study); USES (Uses)  
 (snake venom type I; genotyping methods to detect DNA sequence  
 polymorphisms and haplotypes)

=> d ibib abs hitrn 15

L75 ANSWER 15 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1999:115526 HCPLUS  
 DOCUMENT NUMBER: 130:292382  
 TITLE: High sequence fidelity in a non-enzymic DNA  
 autoligation reaction  
 AUTHOR(S): Xu, Yanzheng; Kool, Eric T.  
 CORPORATE SOURCE: Department of Chemistry, University of Rochester,  
 Rochester, NY, 14627, USA  
 SOURCE: Nucleic Acids Research (1999), 27(3), 875-881  
 CODEN: NARHAD; ISSN: 0305-1048  
 PUBLISHER: Oxford University Press  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The success of oligonucleotide ligation assays in probing  
 specific sequences of DNA arises in large part from high enzymic  
 selectivity against base mismatches at the ligation junction. We describe  
 here a study of the effect of mismatches on a new non-enzymic,  
 reagent-free method for ligation of oligonucleotides. In this  
 approach, two oligonucleotides bound at adjacent sites on a  
 complementary strand undergo autoligation by displacement of a 5'-end  
 iodide with a 3'-phosphorothioate group. The data show that  
 this ligation proceeds somewhat more slowly than ligation by T4 ligase,  
 but with substantial discrimination against single base mismatches both at  
 either side of the junction and a few nucleotides away within one of the  
 oligonucleotide binding sites. Selectivities of >100-fold against  
 a single mismatch are obsd. in the latter case. Expts. at varied concns.  
 and temps. are carried out both with the autoligation of two adjacent  
 linear oligonucleotides and with intramol. autoligation to yield  
 circular "padlock" DNAs. Application of optimized conditions to  
 discrimination of an H-ras codon 12 point mutation is demonstrated with a  
 single-stranded short DNA target.

IT 15181-41-6, Phosphorothioate  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); BIOL (Biological study)  
 (autoligation by displacement of a 5'-end iodide with a 3'-  
 phosphorothioate group; high sequence fidelity in a non-enzymic  
 DNA autoligation reaction)

REFERENCE COUNT: 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT.

&gt; d ind 15

L75 ANSWER 15 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN  
 CC 3-4 (Biochemical Genetics)  
 Section cross-reference(s): 6, 9  
 ST nonenzymic DNA autoligation reaction high sequence fidelity  
 IT Codons  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (12, application of optimized conditions to discrimination of an H-ras  
 codon 12 point mutation is demonstrated; high sequence fidelity in a  
 non-enzymic DNA autoligation reaction)  
 IT DNA  
 RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
 process); BSU (Biological study, unclassified); BIOL (Biological study);  
 PROC (Process)  
 (autoligation; high sequence fidelity in a non-enzymic DNA autoligation  
 reaction)  
 IT Mutation  
 (base-mismatching, ligation proceeds more slowly than ligation by T4  
 ligase, but with discrimination against single base mismatches; high  
 sequence fidelity in a non-enzymic DNA autoligation reaction)  
 IT Gene, animal  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (c-Ha-ras, application of optimized conditions to discrimination of an  
 H-ras codon 12 point mutation is demonstrated; high sequence fidelity  
 in a non-enzymic DNA autoligation reaction)  
 IT DNA  
 RL: BOC (Biological occurrence); BSU (Biological study, unclassified);  
 BIOL (Biological study); OCCU (Occurrence)  
 (circular, autoligation of two adjacent linear oligonucleotides  
 and with intramol. autoligation to yield circular "padlock" DNAs; high  
 sequence fidelity in a non-enzymic DNA autoligation reaction)  
 IT Mutation  
 (point, application of optimized conditions to discrimination of an  
 H-ras codon 12 point mutation is demonstrated; high sequence fidelity  
 in a non-enzymic DNA autoligation reaction)  
 IT 15181-41-6, Phosphorothioate  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
 study, unclassified); BIOL (Biological study)  
 (autoligation by displacement of a 5'-end iodide with a 3'-  
 phosphorothioate group; high sequence fidelity in a non-enzymic  
 DNA autoligation reaction)  
 IT 20461-54-5, Iodide, biological studies  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
 (Biological study); PROC (Process)  
 (autoligation by displacement of a 5'-end iodide with a 3'-  
 phosphorothioate group; high sequence fidelity in a non-enzymic  
 DNA autoligation reaction)

&gt; d 1bib abs hitrn 16

L75 ANSWER 16 OF 17 HCAPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1994:623647 HCAPLUS  
 DOCUMENT NUMBER: 121:223647  
 TITLE: Enzymic preparation of single-stranded DNA containing  
 nuclease-resistant modified nucleotides using  
 phosphorothioate-containing primers  
 INVENTOR(S): Nikiforov, Theo; Knapp, Michael R.  
 PATENT ASSIGNEE(S): Molecular Tool, Inc., USA  
 SOURCE: PCT Int. Appl., 57 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 9  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9416090	A1	19940721	WO 1994-US771	19940118
	W:	AT, AU, BB, BG, BR, BY, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, VN.		
	RW:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG		
US 5518900	A	19960521	US 1993-155746	19931123
AU 9461262	A1	19940815	AU 1994-61262	19940118

AU 674211 B2 19961212  
 EP 679190 A1 19951102 EP 1994-907855 19940118  
 EP 679190 B1 20030502  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE  
 JP 08505535 T2 19960618 JP 1994-516386 19940118  
 JP 3330946 B2 20021007  
 AT 239090 E 20030515 AT 1994-907855 19940118  
 PRIORITY APPLN. INFO.: US 1993-5061 A 19930115  
 US 1993-155746 A 19931123  
 WO 1994-US771 W 19940118

AB A method for generating single-stranded nucleic acid mols. that contain nucleic acid modified nucleotides and so are resistant to 5'-fwdarw.3'-exonucleases are described. The method involves synthesizing the nucleic acid by primer extension using phosphorothioate -contg. primers. A pair of primers with one of them having a phosphorothioate-rich 5'-region and the other not contg. phosphorothioate nucleotides are used to amplify the target sequence. The amplification products are then digested with a 5'-fwdarw.3'-nuclease with the hydrolysis of all of the nucleic acids present except for the amplification products contg. the phosphorothioate-rich primer. These products can be used in DNA sequencing and in the detn. of genetic polymorphism, esp. single base polymorphisms. If the phosphorothioates are placed at the 3'-end of the primer, then any residual primers in the reaction can be hydrolyzed with a 5'-fwdarw.3'-nuclease to prevent further amplification.

=> d ind.16

L75 ANSWER 16 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 IC ICM C12P019-34  
 CC 3-1 (Biochemical Genetics)  
 ST nuclease resistant single stranded DNA  
 IT Deoxyribonucleic acid sequence determination  
 Polymerase chain reaction  
 (enzymic prepn. of single-stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate-contg. primers)  
 IT Genetic polymorphism  
 (single base, detn. of; enzymic prepn. of single-stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate -contg. primers)  
 IT Deoxyribonucleic acids  
 RL: BUU (Biological use, unclassified); RCT (Reactant); BIOL (Biological study); RACT (Reactant or reagent); USES (Uses)  
 (single-stranded; enzymic prepn. of single -stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate-contg. primers)  
 IT Nucleotides, biological studies  
 RL: BUU (Biological use, unclassified); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (oligo-, deoxyribo-, thiophosphate-linked, primers; enzymic prepn. of single-stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate -contg. primers)  
 IT Deoxyribonucleic acids  
 RL: BPN (Biosynthetic preparation); BIOL (Biological study); PREP (Preparation)  
 (thiophosphate-linked, single-stranded, nuclease resistant; enzymic prepn. of single-stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate-contg. primers)  
 IT 79121-99-6, 5'-fwdarw.3'-Exonuclease  
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study).  
 (phage T6 or lambda; enzymic prepn. of single-stranded DNA contg. nuclease-resistant modified nucleotides using phosphorothioate-contg. primers)

=> d ibib abs hitrn 17

✓ L75 ANSWER 17 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN  
 ACCESSION NUMBER: 1992:229075 HCPLUS  
 DOCUMENT NUMBER: 116:229075  
 TITLE: Phosphorothioate-based site-directed mutagenesis for single-stranded vectors  
 AUTHOR(S): Sayers, Jon R.; Eckstein, Fritz

CORPORATE SOURCE: Abt. Chem., Max Planck Inst. Exp. Med., Heidelberg, D-6900/1, Germany

SOURCE: Directed Mutagen. (1991), 49-69. Editor(s): McPherson, M. J. IRL: Oxford, UK.

CODEN: 57RUAL

DOCUMENT TYPE: Conference; General Review

LANGUAGE: English

AB A review with 22 refs. The phosphorothioate-based oligonucleotide-directed mutagenesis method is based on the observation that certain restriction endonucleases are incapable of hydrolyzing phosphorothioate internucleotidic linkages. Thus, double-stranded DNA contg. phosphorothioate linkages in one strand only may be nicked in the non-substituted strand. In this mutagenesis procedure the mismatch oligonucleotide primer is annealed to the (+)strand of a single-stranded circular phage DNA. The primer is extended by a polymn. reaction in which one of the natural deoxynucleoside triphosphates is replaced by the corresponding deoxynucleotide 5'-O-(1-thiotriphosphate), dNTP.alpha.S. Thus, phosphorothioate groups are incorporated exclusively into the (-)strand of the newly synthesized RF-IV DNA. This results in a strand asymmetry which may be exploited. The methods, scope, and limitations of the procedure are discussed.

IT 15181-41-6, Phosphorothioate

RL: BIOL (Biological study)  
(for site-directed mutagenesis of single-stranded DNA vectors)

=> d ind 17

L75 ANSWER 17 OF 17 HCPLUS COPYRIGHT 2003 ACS on STN

CC 3-0 (Biochemical Genetics)

Section cross-reference(s): 9

ST mutagenesis site directed phosphorothioate review

IT Genetic vectors

(single-stranded DNA, site-directed phosphorothioate-based mutagenesis of)

IT Mutation

(site-specific, phosphorothioate-based, for single-stranded DNA vectors)

IT 15181-41-6, Phosphorothioate

RL: BIOL (Biological study)  
(for site-directed mutagenesis of single-stranded DNA vectors)